arXiv:1301.2204v1 [hep-ph] 10 Jan 2013

# Highlights of Supersymmetric Hypercharge ±1 Triplets

Zhaofeng Kang,[1,*] Yandong Liu,[2,†] and Guo-Zhu Ning[1,‡]

[1] *Center for High-Energy Physics, Peking University, Beijing, 100871, P. R. China*

[2] *State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,*
*Chinese Academy of Sciences, Beijing 100190, P. R. China*

(Dated: January 11, 2013)

## Abstract

The discovery of a SM-like Higgs boson with a relatively heavy mass $m_h$ and hints of di-photon excess has deep implication to supersymmetric standard models (SSMs). We consider SSMs extended with hypercharge ±1 triplets, and investigate two scenarios: (A) Triplets couple to the Higgs doublets with significant strength, which can significantly enhance $m_h$ and simultaneously raise the Higgs decay to di-photon rate via the light chargino loops; (B) Oppositely, those couplings are quite weak, but the doubly-charged Higgs bosons, owing to the gauge group structure, offer a natural interpretation to an excess rather than a deficient. Additionally, the pseudo Dirac triplet fermion in this scenario is a natural inelastic non-thermal dark matter candidate. Light doubly-charged particles, especially a robust doubly-charged Higgs boson below about 100 GeV in scenario B, are predicted. We give a preliminary discussion on their search at the LHC.

PACS numbers: 12.60.Jv, 14.70.Pw, 95.35.+d

---

[*] Electronic address: `zhfkang@gmail.com`

[†] Electronic address: `ydliu@itp.ac.cn`

[‡] Electronic address: `ngz@gmail.com`

## I. INTRODUCTION AND MOTIVATIONS

The LHC is on the verge of claiming the discovery of a standard model (SM)-like Higgs boson [1] (Now the significance reaches 6.9 $\sigma$). This good news consolidates supersymmetry (SUSY), which elegantly solves the gauge hierarchy problem relating with this putatively fundamental spin-0 particle, as the leading candidate for new physics. However, we encounter two problems. Firstly, the measured SM-like Higgs boson mass $m_h$ is about 126 GeV, a sufficiently large value to seriously challenge the naturalness of the weak scale in the most popular minimal supersymmetric SM (MSSM) [2]. Secondly, in terms of the announced results [1], this Higgs boson is not exactly SM-like and its di-photon signature channel shows the most noticeable deviation, to which the MSSM also lacks a natural explanation. Thus, solving these two problems (in spite of the waiting-for-confirmation for the latter) simultaneously and naturally gives us illuminating hints to go beyond the MSSM.

Actually, the relatively heavy Higgs boson mass alone may open a window for new models. As been well known, in the MSSM the tree-level SM-like Higgs boson mass is upper bounded by the $Z-$boson mass, owing to the fact that the strength of Higgs quartic term is determined by the electroweak gauge couplings. The new models should provide significant Higgs quartic terms either at the tree- or loop-level. A well known example is the singlet extended MSSM such as the next-to MSSM (NMSSM) which possesses a large coupling between the singlet and Higgs doublets. Additionally, it shows a singlet-doublet mixing effect to enhance $m_h$ [2, 3] (The relat4ed search please see Ref. [4]). Recently, some other exploration beyond the MSSM alone this line includes: (A) The gauge group extension, which gives non-decoupling $D-$terms [5] to increase $m_h$ (It requires the Higgs doublets to be effectively charged under the gauge group); (B) Vector-like particles extension, which raise the Higgs boson mass at the loop-level [6]; (C) $SU(2)_L$ triplet $T_0$ with hypercharge 0 extension [7, 8]. $T_0$ couples to the Higgs doublets via $\lambda_{T_0}\mathrm{Tr}H_uT_0H_d$ which, in enhancing $m_h$, shows similar behavior to the NMSSM.

However, once we further impose the global fit for the Higgs data [9], they are difficult to provide a sufficiently good interpretation. The global fit needs new QED charged particles which can directly enhance the width of Higgs decay to di-photon [9] without significantly effecting other Higgs tree-level couplings, especially to $ZZ/WW$. Within the NMSSM, light charginos, prediction of natural SUSY, may be the viable charged particle candidate in the case of a large singlet-doublet mixing [10]. But this scenario heavily depends on the tuned mixing structure. In the case of an enlarged presentation such as $T_0$, new charged fermions are introduced, and therefore they can lead to larger di-photon rate given a large enough $\lambda_{T_0}$ [8].

In this paper we consider the low energy SUSY incorporating $SU(2)_L$ triplets which carry hypercharge ±1, denoted as $T_{u,d}$ respectively. As been noticed long ago by [11] and more recently [12–14], such fields can effectively lift $m_h$ if they couple to the Higgs doublets with significant couplings. Even more interestingly, they, the singly-charged charginos mixed with the MSSM partners, are capable of increasing the di-photon rate simultaneously. Thereby, such a SSM is a good case following the rule at the beginning. Driven by the doubly-charged Higgs bosons, the $T_{u,d}$ weak Yukawa coupling limit is also explored. This scenario presents an elegant explanation to why the doubly-charged Higgs boson-loop constructively rather than destructively interfere with the $W-$loop. Additionally, this scenario provides a pseudo Dirac fermion to be an inelastic non-thermal dark matter (DM) candidate. Finally, both scenarios predict quite light charged states and in particular, the doubly-charged charginos and Higgs (for the latter scenario only) may have a good LHC discovery prospect.

This paper is organized as follows. In Section II we study the MSSM extended with hypercharge ±1 triplets, two distinct scenarios are explored and we stress their phenomenological highlights for Higgs data and dark matter. Corresponding LHC search strategy and prospect are also briefly commented. Discussion and conclusion are casted in Section III and some necessary and complementary details are given in the Appendices.

## II. HIGHLIGHTS OF SUSY WITH HYPERCHARGE ±1 TRIPLETS

### A. Model setup

Naturalness is an important guidance to explore the structure of the low energy SSMs. In this section, motivated by naturally explaining the recent Higgs data, including the relatively heavy SM-like Higgs boson mass and the hinted excess of Higgs decay to di-photon rate, we consider $SU(2)_L$ triplets $T_{u,d}$ extended SSMs (TSSMs). Here $T_{u,d}$ carry hypercharges $-1$ and $+1$, respectively. Note that both of them are needed for the sake of anomaly cancelation. The most relevant modula of the TSSM is

$$W_H = \mu H_u \cdot H_d + \mu_T \mathrm{Tr}(T_u T_d) + \lambda_u H_u \cdot T_u H_u + \lambda_d H_d \cdot T_d H_d. \tag{1}$$

The corresponding soft SUSY-breaking Lagrangian is

$$\begin{aligned}\mathcal{L}_{soft} = & \sum_{\Phi=H_{u,d},T_{u,d}} |\Phi|^2 + (B\mu H_u \cdot H_d + B\mu_T \mathrm{Tr}(T_u T_d) + c.c.) \\ & + (A_u H_u \cdot T_u H_u + A_d H_d \cdot T_d H_d + c.c.)\,. \end{aligned} \tag{2}$$

For simplicity, here all parameters are assumed to be real (Incorporating CP violation will bring about quite different Higgs phenomenologies [15].). $T_u$ and $T_d$, following the notation of Ref. [11, 13], are respectively written as

$$T_u \equiv T_u^a \sigma^a = \begin{pmatrix} \frac{T_u^-}{\sqrt{2}} & -T_u^0 \\ T_u^{--} & -\frac{T_u^-}{\sqrt{2}} \end{pmatrix}, \quad T_d \equiv T_d^a \sigma^a = \begin{pmatrix} \frac{T_d^+}{\sqrt{2}} & -T_d^{++} \\ T_d^0 & -\frac{T_d^+}{\sqrt{2}} \end{pmatrix}, \tag{3}$$

with $\sigma^a$ the three Pauli matrices. As usual, the doublets are written as $H_u = (H_u^+, H_u^0)$ and $H_d = (H_d^0, H_d^-)$. For illustration, in components the superpotential terms take the form of

$$H_u \cdot T_u H_u \equiv (Hu)_\alpha \epsilon^{\alpha\beta} (T_u)_\beta{}^\gamma (H_u)_\gamma = \sqrt{2} H_u^+ H_u^0 T_u^- - (H_u^0)^2 T_u^0 - (H_u^+)^2 T_u^{--}, \tag{4}$$

where the antisymmetric tensor $\epsilon^{\alpha\beta}$ has entry $\epsilon^{12} = -1$. The above definition of spinor product is applied to other quantities.

If we want to produce the small neutrino masses, as in the type-II seesaw mechanism [16], the following lepton number violating operators are introduced:

$$W_L = (\lambda_L)_{ij} L_i \cdot T_d L_j. \tag{5}$$

with $i, j = 1, 2, 3$ the family indices. Through the products of the small Yukawa couplings and triplet's vacuum expected value (VEV), they generate the tiny neutrino masses $(m_\nu)_{ij} = (\lambda_L)_{ij} v_{T_d} \sim 10^{-10}$ GeV. One will find, the triplets' VEV of interest should be around the GeV scale, as renders the Yukawa couplings extremely small. Then, alternatively, one may forbid such couplings and turn to the canonical seesaw mechanism. In any case, this aspect of the TSSM is not of our main concern. But we would like to stress that, models under consideration can be embedded into the supersymmetric type-II seesaw mechanism [17] without violating the $R-$parity, namely, the lightest sparticle LSP DM hypothesis of SUSY can be maintained. In contrast, the models with hypercharge 0 triplets can not be embedded into the type-III seesaw mechanism and meantime respect the $R-$parity.

In the TSSM, $\lambda_u$ and $\lambda_d$ are closely related to the ensuing discussions. In the one scenario studied later they will be required to be large, so we want to know the scale at which the perturbitivity breaks. The relevant renormalization group equations (RGEs) can be found in Appendix. A. From it one can find, due to $h_t \gg h_b$, the beta functions of $\lambda_u$ and $\lambda_d$ are asymmetric. $\lambda_d$ is favored to be moderately larger than $\lambda_u$. Fig. 1 shows that if they do not hit the Landau pole below the GUT-scale, the maximal $\lambda_d$ and $\lambda_u$ are about 0.62 and 0.45, respectively. While for $\lambda_d = 0.7$, it hits the Landau pole soon at a scale $\sim 10^{10}$ GeV. The implications to GUT is beyond the scope of this work and we simply discuss it in the end of the work.

In the following subsections we will turn our attention to the highlights for the TSSMs, focusing on phenomenologies involving the Higgs and DM. Two different scenarios are investigated separately.



FIG. 1: The plot of $\lambda_{u,d}$ verses the running scale. Two groups of boundary values at the low energy are chosen: (A) $\lambda_u = 0.3$ (thick dashed line) and $\lambda_d = 0.7$ (thick line); (B) $\lambda_u = 0.45$ (thin dashed line) and $\lambda_d = 0.62$ (thin line), showing the maximal $\lambda_{u,d}$ endured by perturbativity up to the GUT-scale.

### B. Scenario A: Large $\lambda_{u,d} \sim 1$

This scenario takes the advantage in rasing the SM-like Higgs boson mass and at the same time the Higgs decay width to di-photon. But these two aspects tend to show a tension and the mixing effect can relax it.

#### *1. Lifting the Higgs boson mass and pull-down mixing effect*

We now study the CP-even Higgs boson sector, which consists of four states, with two from the ordinary doublets and other two from the triplets $T^0_{u,d}$. The triplets are expected to nearly decouple from the doublets because their mixings are induced by the triplets' VEV, which however should be no more than a few GeVs otherwise the $\rho$ parameter will deviate from 1 beyond the experimentally allowed level [13]. In fact, the triplets' VEVs can be approximately determined in an analytical way:

$$\begin{aligned} v_{T_{u,d}} \equiv& \langle T^0_{u,d} \rangle \simeq \frac{v^2}{m^2_{T_{u,d}} + \mu_T^2} \mathcal{M}_{u,d}, \\ \mathcal{M}_u \equiv& A_u \sin^2\beta - \lambda_d \mu_T \cos^2\beta - \lambda_u \mu \sin 2\beta, \\ \mathcal{M}_d \equiv& A_d \cos^2\beta - \lambda_u \mu_T \sin^2\beta - \lambda_d \mu \sin 2\beta, \end{aligned} \tag{6}$$

with $\tan\beta = \langle H^0_u \rangle / \langle H^0_d \rangle$. The approximations are valid in the heavy scalar triplets limit and moreover a not anomalously large $B_{\mu_T}$. As one can see, the smallness of $\langle T^0_{u,d} \rangle \lesssim 1$ GeV, in the case of a light $\mu_T \sim 100$ GeV while large $\lambda_{u,d} \sim 1$, can be feasible only if $m_{T_{u,d}} \gtrsim \mathcal{O}(1)$ TeV. Otherwise, we have to greatly tune parameters to make $\mathcal{M}_{u,d}$ very small. On the other hand, the triplet $T_u$ couples to $H_u$ with a large Yukawa coupling, so $m^2_{H_u}$ will receive a large negative correction via the renomalization group equation (RGE):

$$16\pi^2 \frac{dm^2_{H_u}}{dt} = 6h_t^2 \left(m^2_{Q_3} + m^2_{\tilde{U}^c}\right) + 12\lambda_u^2 m^2_{T_u} + ..., \tag{7}$$

In the leading logarithm and allowing for a fine-tuning $F = 100$, we obtain an upper bound

$$m_{T_u} \lesssim 1.2 \times \left(\frac{F^{1/2}}{10}\right)\left(\frac{0.5}{\lambda_u}\right)\left(\frac{30}{\log \Lambda/\text{TeV}}\right)^{1/2} \text{TeV}, \tag{8}$$

where $\Lambda$ is the SUSY-breaking mediation scale and in the above estimation we have set it to be the grand unification theory (GUT)-scale. Even if we set $\Lambda = 10^6$ GeV, the upper bound is merely doubled. In other words, generically speaking the VEVs can not lie far below one GeV. This has important implication to the mixing effect.

To further quantitatively understand the above arguments, here we have a close inspection into the mass square matrix. We adopt the following field decomposition

$$\begin{aligned} H_u^0 =& v_u + \frac{1}{\sqrt{2}}(h_1 \cos\beta + h_2 \sin\beta) + \frac{i}{\sqrt{2}}(P_1 \cos\beta + G^0 \sin\beta), \\ H_d^0 =& v_d + \frac{1}{\sqrt{2}}(-h_1 \sin\beta + h_2 \cos\beta) + \frac{i}{\sqrt{2}}(P_1 \sin\beta - G^0 \cos\beta), \\ T_u^0 =& v_{T_u} + \frac{1}{\sqrt{2}}(h_3 + iP_2), \quad T_d^0 = v_{T_d} + \frac{1}{\sqrt{2}}(h_4 + iP_3). \end{aligned} \tag{9}$$

Then, in the basis $(h_1, h_2, h_3, h_4)$ the mass square matrix $M_S^2$ is given by Eq. (B3). In particular, the SM-like Higgs boson mass square, ignoring both the doublet-doublet and doublet-triplet mixing effects, is given by:

$$(M_S^2)_{22} = m_Z^2 \left[\cos^2 2\beta + \frac{4}{g^2}\left(\lambda_d^2 \cos^4\beta + \lambda_u^2 \sin^4\beta\right)\right], \tag{10}$$

with $m_Z^2 = g^2 v^2$ and $g^2 = (g_1^2 + g_2^2)/2$. The new contributions originate from $\lambda_{u.d} H_{u.d} T_{u.d} H_{u.d}$, which give new quartic terms such as $|F_{T_u^0}| \supset \lambda_u^2 |H_u^0|^4$. So, provided that order one couplings $\lambda_{u,d}$, the tree-level Higgs boson mass can be sufficiently large regardless of $\tan\beta$.

Such an interesting feature, obviously, is attributed to the fact that the quartic terms $|H_d^4|$ and $|H_u^4|$ are generated simultaneously. To our knowledge, basically the $m_h$ dependence on the angle $\beta$ can be classified into three types: (I) The MSSM-like case where the quartic term is determined by the vector-like $D-$terms (Namely $H_{u,d}$ form a vector-like expression under the gauge group, such as $SU(2)_L \times U(1)_Y$), then the quartic term $(|H_u^0| - |H_d^0|)^2 \propto \cos^2 2\beta$ which requires a large $\tan\beta$; (II) The NMSSM-like case, the quartic term is a two-doublets mixed pattern $|H_u^0 H_d^0|^2 \propto \sin^2 2\beta$ and thus a small $\tan\beta \sim 1$ is required. The SSMs with hypercharge-0 triplets [7, 8] also fall into this category; (III) The TSSMs considered in this work as well as in models extended by a new gauge group under which $H_u$ and $H_d$ form a chiral presentation. Such a classification is useful, since the di-photon excess usually also has a strong dependence on $\beta$, as will be explicitly seen soon. Therefore it provides a helpful guild to actual model building.

The maximal enhancement indicated in Eq. (10), however will be reduced by the mixing effect. As suggested previously, decoupling of the triplets is merely marginal when we set the rough naturalness bound $m_{T_{u,d}} \lesssim 2$ TeV (In reality, $T_d$ can be much heavier since it does not couple to $H_u$, but we do not consider such a hierarchy.). To see that, we start from a simplified system by only including the sub-matrix consisting of entries involving states 2 and 3 (namely only including the mixing with $T_u^0$), and from the expressions given in Eq. (B3) it is not difficult to find the relation

$$(M_S^2)_{22}(M_S^2)_{33} \sim \mathcal{O}(0.1)(M_S^2)_{23}^2. \tag{11}$$

Thereby the mixing effect will pull-down the lighter eigenvalue (namely the SM-like Higgs boson mass square) by a mount $(M_S^2)_{23}^4/(M_S^2)_{33}$ [2] and renders a reduce to the Higgs boson mass, with a amount estimated by

$$\begin{aligned}\Delta m_h \approx& -\frac{1}{2}(M_S)_{22}\frac{(M_S^2)_{23}^4}{(M_S^2)_{33}(M_S^2)_{22}}\\ =& -2\left(\frac{\mathcal{M}_u^2}{m_{T_u}^2}\right)\left(\frac{m_h}{(M_S)_{22}}\right)\left(\frac{v^2}{m_h}\right)\text{ GeV}.\end{aligned} \tag{12}$$

The mixing with $T_d^0$ similarly contributes to the pulling-down effect and then the estimation should be doubled. Through mild tuning one can obtain a relatively small $\mathcal{M}_{u,d}$. It helps to not only reduce $v_{T_{u,d}}$, which is consistent with the neutrino phenomenology and the $\rho$ parameter constraint, but also weaken the pulling-down effect. However, sometimes the new quartic terms excessively enhance $m_h$, as is just the case when we want to further account for the di-photon excess, then we need a relatively large $\mathcal{M}_{u,d}$, or equivalently $v_{T_{u,d}}$, to strength the pulling-down effect. From Eq. (6) we know a negative $A_{u,d}$ are favored then. We plot the two limits of mixing effect on Fig. 2.

### *2. Rasing the Higgs di-photon rate via charginos*

The rate of Higgs decay to di-photon is sensitive to charged particles beyond the SM. Given such particles, one can predict its signature strength, which is defined as the ratio of



FIG. 2: Contour plots of $m_{T_u}$ (solid lines) and the amount of reducing $m_h$ (dashed lines) from the doublet-triplet pulling-down mixing effect on the $v_{T_u} - A_u$ plane. Left: Significant pulling-down with $\lambda_d = 0.70$, $\lambda_u = 0.45$, $v_{T_d} = -2.0$ GeV and $A_d = -300$ GeV. Right: Weaker pulling-down by setting $\lambda_d = 0.70$, $\lambda_u = 0.30$, $v_{T_d} = -0.5$ GeV and $A_d = 200$ GeV. Other not much relevant parameters are common to both cases: $\mu = 200$ GeV, $\mu_T = 250$ GeV, $B\mu = 400 \times 200$ GeV$^2$, $B\mu_T = 200^2$ GeV$^2$ and $\tan\beta = 1.5$ (Actually, effect of $\mu$, $\mu_T$ and $\tan\beta$ can be absorbed into $A_{u,d}$, see Eq. (6)).

the di-photon rates

$$R_{\gamma\gamma} \equiv \frac{\sigma(gg \to h)\mathrm{Br}(h \to \gamma\gamma)}{\sigma_{\mathrm{SM}}(gg \to h)\mathrm{Br}_{\mathrm{SM}}(h \to \gamma\gamma)}. \tag{13}$$

In general, the branching width of Higgs decay to di-photon is formulated as [18]

$$\Gamma(h \to \gamma\gamma) = \frac{\alpha^2 m_h^3}{1024\pi^3} \left| \frac{g_{hVV}}{m_V^2} Q_V^2 \mathcal{A}_1(\tau_V) + \frac{2g_{hf\bar{f}}}{m_f} N_{c,f} Q_f^2 \mathcal{A}_{1/2}(\tau_f) + \frac{g_{h|S|^2}}{m_S^2} N_{c,s} Q_S^2 \mathcal{A}_0(\tau_S) \right|^2, \tag{14}$$

with $N_c$ the color factor and $\tau_i \equiv 4m_i^2/m_h^2$. Here $V$, $f$, and $S$ denote for a vector boson, Dirac fermion and charged scalar, respectively. Their charges and couplings to the Higgs boson are labeled as $Q_{V,f,S}$ and $g_{hVV}$, etc. In the limit $\tau_i \gg 1$, which is hold in this work, the loop-functions take the asymptotic values $\mathcal{A}_1 \to -7$, $\mathcal{A}_{1/2} \to 4/3$ and $\mathcal{A}_0 \to 1/3$. Specified to the SM, the $W-$ and top-loop dominantly contribute to this width and give

$$\mathcal{A}_1(\tau_W) \approx -8.32, \quad 3 \times (2/3)^2 \mathcal{A}_{1/2}(\tau_t) \approx 1.84, \tag{15}$$

where $m_h = 126$ GeV is fixed. In the supersymmetric models extended by the triplets $T_{u,d}$ with hypercharge $\pm 1$, the singly-charged charginos and doubly-charged Higgs bosons are two promising candidates to make $R_{\gamma\gamma} > 1$. But the doubly-charged charginos, in spite of their larger charges, are irrelevant because of their absence of direct coupling with the neutral Higgs bosons. This can be explicitly seen from the superpotential.

We now consider the contributions from the light charginos. In the first place, for a mixed Driac system $f_i$ with a mass matrix $M_F$ (Its eigenvalues are properly large), the coupling $\sum_i 2g_{hf_i\bar{f}_i}/m_{f_i}$ (with $f_i$ in the mass eigenstate) can be obtained via [18, 19]

$$\sum_i \frac{2g_{hf_i\bar{f}_i}}{m_{f_i}} = \frac{1}{\sqrt{2}}\frac{\partial}{\partial v}\log\left(\det M_F^\dagger M_F\right). \tag{16}$$

Applied to the top quark, it gives $g_{ht\bar{t}} = \sqrt{2}/v$ with $v = 174$ GeV. In this paper, the chargino system consists of three pairs, from winos $\widetilde{W}^\pm$, Higgsinos $\widetilde{H}^\pm_{u,d}$ as well as the triplets $\widetilde{T}^\pm_{u,d}$. The chargino mass matrix is given by

$$(-i\widetilde{W}^-, \widetilde{H}^-_d, \widetilde{T}^-_u)\begin{pmatrix} M_2 & g_2 v_u & 0 \\ g_2 v_d & \mu & \sqrt{2}\lambda_d v_d \\ 0 & \sqrt{2}\lambda_u v_u & \mu_T \end{pmatrix}\begin{pmatrix} -i\widetilde{W}^+ \\ \widetilde{H}^+_u \\ \widetilde{T}^+_d \end{pmatrix}, \tag{17}$$

where terms proportional to $v_{T_{u,d}}$ are safely dropped. Then the sum of the couplings between $h$ and $\bar{\chi}_i\chi_i$, with $\chi_i\,(i = 1,2,3)$ the three singly-charged charginos in the mass eigenstates, is calculated to be

$$\frac{2g_{h\bar{\chi}\chi}}{m_\chi} \equiv \sum_{i=1}^{3}\frac{2g_{h\bar{\chi}_i\chi_i}}{m_{\chi_i}} = \frac{2\sin 2\beta\,(2\lambda_d\lambda_u v/\mu_T + g_2^2 v/M_2)}{\sin 2\beta\,(2\lambda_d\lambda_u v/\mu_T + g_2^2 v/M_2) - 2\mu/v}\frac{\sqrt{2}}{v}. \tag{18}$$

Immediately, it is seen that a small $\tan\beta \sim 1$, as in the MSSM, is an indispensable condition [35]. Furthermore, compared to the MSSM case, our result in fact can be obtained just through a shift: $g_2^2 v/M_2 \to g_2^2 v/M_2 + 2\lambda_d\lambda_u v/\mu_T$. So either $\lambda_u\lambda_d \ll 1$ or $\mu_T \gg v$, the triplets will decouple at the loop generating the Higgs di-photon decay. Under such limits the result is reduced to the MSSM case, from which the di-photon enhancement is no more than 30% even we work in the unrealistic limit with $\tan\beta = 1$ [20].

So, the non-decoupling triplets case with properly light $\widetilde{T}^\pm_{u,d}$ and moreover $\lambda_d\lambda_u \sim \mathcal{O}(1)$ is of interest. It is, in spite of some tension, consisting with raising the Higgs boson mass. Now the Higgs di-photon rate signature strength is given by

$$R_{\gamma\gamma} \approx \left|1 - \frac{1.84}{8.33 - 1.84} \times \frac{3}{4}\frac{v}{\sqrt{2}}\frac{2g_{h\bar{\chi}\chi}}{m_\chi}\right|^2, \tag{19}$$

where we have used the result Eq. (15) and normalized the new physics contribution to the top quark contribution, indicated by the second term above. Moreover, we assume that all the charginos' loop functions $\mathcal{A}_{1/2}(\tau_{\chi_i})$ are equal to $\mathcal{A}_{1/2}(\tau_t)$. Such an approximation typically underestimates the contribution since the lightest chargino mass is about half of the top quark mass, which leads to a slight enhancement in the loop function compared to the one of top quark.

Numerically results, combined with the constraint from Higgs boson mass, are shown on the left panel of Fig. 3. From it one can find that: (I) All the charginos, including the winos, should be properly light. The fact that the light winos are favored is not surprising, and it just indicates that the MSSM charginos should have a non-negligible contribution to $R_{\gamma\gamma}$, This is important to get a large $R_{\gamma\gamma}$ without turning to a too large $\lambda_u\lambda_d$; (II) Large di-photon rate may have a tension with $m_h = 126$ GeV, but then the previous pulling-down effect can relax it; (III) Note that only the product $\lambda_u\lambda_d$ is relevant to determine $R_{\gamma\gamma}$, so, in light of the discussion in Section II A, we can take $\lambda_u = 0.4$ and $\lambda_d = 0.62$ to get viable enhancements without spoiling perturbativity below the GUT-scale. Finally, the sign of $g_{h\bar{\chi}\chi}$, as in most models, is a result of parameter tuning (or artificial choice). In the following, we will present an attractive scenario to naturally understand the origin of the sign.

### C. Scenario B: The small $\lambda_{u,d} \ll 1$ limit

We now switch to the scenario characterized by very small couplings $\lambda_{u,d} \ll 1$, which can not raise $m_h$ but is still of great interest. In the light of Eq. (6), one can find that the doubly-charged Higgs bosons now can be as light as the weak-scale particles without spoiling the $\rho$ parameter. Then, although the charginos' contribution is reduced to the MSSM level, instead $H^{\pm\pm}$ potentially lead to a considerable $R_{\gamma\gamma}$ enhancement. Actually, in the following we will show that it provides a very interesting way to understand this enhancement and moreover is predictive. On top of that, this scenario naturally offers an inelastic dark matter (DM) candidate.

#### 1. *Rasing the di-photon rate via doubly-charged Higgs bosons*

Although the superpotential does not provides coupling between $H^{\pm\pm}$ and the SM-like Higgs boson, $D-$term does. The interesting point is that, by virtue of supersymmetry, those couplings are fixed by the model gauge structure rather than input by hand. Explicitly, from



FIG. 3: Left: A contour plot of di-photon excess strength on the $\mu_T-\lambda_u$ plane. For illustration, two cases are demonstrated: (A) $\lambda_d=0.7$, $\mu=200$ GeV, $M_2=300$ GeV and $\tan\beta=1.5$. The direct charigno lower bound $m_{\chi_1}>94$ GeV [22] (thick red line) is imposed. The two thick black lines label $R_{\gamma\gamma}=1.3$ and 1.5 respectively. Additionally, we allow Higgs boson mass moderately heavier than 125 GeV, indicated by the green lines, by virtue of the possible pulling-down effect to reduce them, which is discussed in the text. The region filled by blue dashed lines denote the viable window for this case; (B) The only difference is here we take $M_2=600$ GeV, and various bounds are labeled by dashed lines. The viable window, filled by red points, is significantly narrower than case A. Right: A plot of di-photon excess $R_{\gamma\gamma}$ from the doubly-charged Higgs boson varying with its mass (10% reduce from the singly-charged Higgs boson has been included.). Regardless of any other model details, an enhancement about $R_{\gamma\gamma}\sim 1.3$ predicts a doubly-charged Higgs boson around 90 GeV.

Eq. (B1) it is not difficult to extract the following couplings

$$\begin{aligned}
V_D \supset & \frac{g_2^2-g_1^2}{2}(|H_u^0|^2-|H_d^0|^2)\left(|T_u^{--}|^2-|T_d^{++}|^2\right)\\
& -\frac{g_1^2}{2}(|H_u^0|^2-|H_d^0|^2)\left(|T_u^{-}|^2-|T_d^{+}|^2\right)\\
= & -\sqrt{2}\cos 2\beta\cos 2\theta_w\frac{m_Z^2}{v}h_2\left(|T_u^{--}|^2-|T_d^{++}|^2\right)\\
& +\sqrt{2}\cos 2\beta(\sin^2\theta_w/\cos 2\theta_w)\frac{m_Z^2}{v}h_2\left(|T_u^{-}|^2-|T_d^{+}|^2\right),
\end{aligned}\tag{20}$$

where $\theta_w$ is the Weinberg angle and $\sin^2\theta_w\approx 0.23$. As a consequence of $T_u^{--}$ lying at the 21 position in $T_u$ while $T_d^{++}$ at the 12 position in $T_d$, there is a relative sign between the two terms in Eq. (20) (Essentially, the reason can be traced back to the hypercharge assignment.). Therefore, if $T_u^{--}$ is very heavy, say due to a large $m_{T_u}^2 + \mu_T^2 \gg m_{T_d}^2 + \mu_T^2$ (Such kind of hierarchy can be naturally realized in the gauged mediated SUSY-breaking models by coupling $T_u$ to messengers [21].), then the second term dominates. Similarly, we obtain the coupling as Eq. (16):

$$\frac{g_{h|T_d^{++}|^2}}{m_{T_d^{++}}^2} = \cos 2\beta \cos 2\theta_w \frac{m_Z^2}{m_{T_d^{++}}^2} \frac{\sqrt{2}}{v}. \tag{21}$$

The singly-charged Higgs boson with comparable mass reduces the doubly-charged Higgs boson contribution $\approx 10\%$ and will be simply took into account hereafter. It is noticed that Eq. (21) takes a negative sign which, alone with its strength, is totally determined by the gauge structure except for the dependence on mass.

The strength of enhancement $R_{\gamma\gamma}$ is significant only in the large $\tan\beta$ limit. Moreover it sensitively depends on the doubly-charged Higgs boson mass, and an appreciable amount implies that the mass should be no heavier than 100 GeV. Concretely, we now have

$$R_{\gamma\gamma} \approx \left| 1 + \frac{1.84}{8.33 - 1.84} \times \frac{m_Z^2}{m_{T_d^{++}}^2} \frac{3\mathcal{A}_0(\tau_{T_d^{++}})}{\mathcal{A}_{1/2}(\tau_t)} \right|^2. \tag{22}$$

It has a sole parameter $m_{T_d^{++}}$ and the left panel of Fig. 3 shows that an excess $R_{\gamma\gamma} \gtrsim 20\%$ clearly demands a doubly-charged Higgs boson with mass below 105 GeV.

Comments are in orders. First, although there is a great number of works using a doubly-charged Higgs field to enhance the di-photon rate [23], here we find that supersymmetry gives the most elegant realization. Second, from the above deduction it is tempting to conjecture that, this sign-determination mechanism may be applied to any supersymmetric models, which contain at least one gauge group $G$ and under it both the Higgs doublets and some QED-charged chiral fields form the vector-like presentation. Third, as mentioned at the beginning, the TSSM in this scenario has to give up the merit of lifting the Higgs boson mass [36]. But this is not a server problem since any mechanism reviewed in the introduction, or simply heavy stops, could provide a solution. But note that the NMSSM-like model is not viable since it requires a small $\tan\beta$.

### 2. *Pseudo-Dirac fermions as a non-thermal inelastic DM candidate*

In the small $\lambda_{u,d}$ scenario, we can show that the model naturally provides an inelastic DM candidate [25, 26]. Within this scenario, owing to the requirement of di-photon enhancement, $\mu_T$, barring large cancelation, is favored to be as small as possible so that ${T_d}^{++}$ is naturally light. Furthermore, the $\rho$ parameter typically requires $\lambda_{u,d} \lesssim 10^{-2}$. These two features of the model parameter space inspire an interesting scenario: The nearly secluded (in the sense of superpotential) fermionic triplets's neutral components $\widetilde{T}^0_{u,d}$ form a Dirac pair with mass $\mu_T$, however, the Dirac pair is slightly split by the mixing with the MSSM neutralinos. As a consequence, the Dirac fermion becomes pseudo Dirac. Concretely, the 6 by 6 neutralino mass matrix, in the basis $\chi_0^T = (-i\widetilde{B}, -i\widetilde{W}^3, \widetilde{H}^0_d, \widetilde{H}^0_u, \widetilde{T}^0_d, \widetilde{T}^0_u)$, takes a form of

$$M_{\chi^0} \approx \begin{pmatrix} M_1 & 0 & -\frac{g_1 v_d}{\sqrt{2}} & \frac{g_1 v_u}{\sqrt{2}} & \sqrt{2}g_1 v_{T_d} & -\sqrt{2}g_1 v_{T_u} \\ & M_2 & \frac{g_2 v_d}{\sqrt{2}} & -\frac{g_2 v_u}{\sqrt{2}} & -\sqrt{2}g_2 v_{T_d} & \sqrt{2}g_2 v_{T_u} \\ & & 0 & -\mu & \frac{v_d \lambda_d}{\sqrt{2}} & 0 \\ & & & 0 & 0 & \frac{v_u \lambda_u}{\sqrt{2}} \\ & & & & 0 & \mu_T \\ & & & & & 0 \end{pmatrix}. \tag{23}$$

The analytical analysis is very difficult, but we can capture the primary information of the pseudo Dirac system, i.e., the mass splitting in an approximate way.

At the leading order, the mass spitting is attributed to the mixing between the Dirac triplets and particles with Majorana mass term, say the wino (The splitting from mixing with bino is assumed to be sub-leading for the momentum, owing to the smaller gauge coupling $g_1 \ll g_2$). In this sense we can simply decouple the Higgsinos (A small $\mu$ tends to reduce the splitting), drop bino and further work in the limit $v_{T_u} \ll v_{T_d}$ (This is ture since $m^2_{T_u} \gg m^2_{T_d}$), and then obtain the greatly simplified three by three matrix

$$M_{\chi^0} \approx \begin{pmatrix} M_2 & -\sqrt{2}g_2 v_{T_d} & 0 \\ -\sqrt{2}g_2 v_{T_d} & 0 & \mu_T \\ 0 & \mu_T & 0 \end{pmatrix}. \tag{24}$$

The mass splitting is calculated to be

$$\delta = \frac{M_2}{M_2^2 - \mu_T^2}(\sqrt{2}g_2 v_{T_d})^2. \tag{25}$$

As one can see, for a sub-GeV triplet VEV and moreover moderately heavy $M_2 \sim 500$ GeV, the splitting is naturally at the sub-MeV scale. Dropping bino renders the above estimation only about half of the actual value, if the bino is considerably light, say $M_1 \simeq M_2/3$ suggested by the gaugino mass unification (Actually, one can include the bino contribution by adding a similar term to Eq. (25) with the replacement $M_2 \to M_1$ and $g_2 \to g_1$). The full result accommodates a split over a rather wide range.

A weak scale inelastic DM with sub-MeV mass splitting has several phenomenologically interesting applications. In the first place, it provides a way to avoid the stringent spin-independent bound on the $Z-$mediated DM-nucleon recoil, which is very important to make the triplets DM model here not been excluded by the DM direct detection experiments [25, 27] like XENON100 [28] (It, along with the WMAP, renders the conventional neutralino LSP DM now not favored by naturalness [2]). Secondly, it is a possible candidate to explain the 511keV line [29], the DAMA/LIBRA experiment result [26], or for some other purposes [30].

However, as been well known, a DM particle participating in full weak gauge interactions will have a too large annihilation cross section $\sigma v \sim g_2^4/16\pi\mu_T^2$, via the weak gauge bosons mediation or directly into them. Consequently, the DM relic density is too small to account for its observed energy fraction in our Universe today. The exception is that the DM mass is at the TeV scale, but that is inconsistent with the scenario considered here. Thus we should go beyond the thermal DM scenario. For instance, the ordinary LSP (OLSP) is highly bino-like (Thus it does not couple to the $Z-$boson), which is assumed to acquire a proper thermal relic density and late decays to the pseduo-Dirac fermion plus a pair of light fermions, mediated by a slepton (See a similar scenario in a different context [24].). The decay width is estimated to be

$$\Gamma(\widetilde{B} \to \widetilde{T}^0_{u,d} f f') \sim \frac{1}{(4\pi)^3} g_1^4 V_{14}^2 \frac{M_1^5}{m^4_{\widetilde{f}}} P(M_1^2/\mu_T^2) \sim 10^{-18}\text{GeV}. \tag{26}$$

In this estimation, we have set the mixing angle between bino and triplet fermion at the order $V_{14} \sim g_1^2 v_{T_d}^2/(\mu_T m_{\widetilde{B}}) \sim 10^{-4}$, a TeV scale slepton as well as the phase space suppressing factor $P \sim 10^{-2}$ which, in principle can be even smaller in magnitude provided sufficient degeneracy between bino and DM. Hence the decay can happen after the DM freezing-out, and the bino can transfer its number density to the DM.

### D. The preliminary of LHC search

As is expected, to sufficiently enhance the di-photon rate, new light charged particles will appear. In the TSSM, in any scenario the fermionic triplets should be rather light while in the scenario B the triplets scalar components must stay light, concretely, the doubly charged Higgs boson mass lying below 100 GeV. As a result, the doubly-charged charginos (denoted as $\chi^{++}$) and Higgs bosons (denoted as $H^{++}$) are likely to be probed at the LHC due to their clear signature, i.e., multi same-sign leptons plus missing energy. In this subsection, as a preliminary discussion, we will briefly comment on their prospect at the near LHC running but leave more quantitative study elsewhere.

### 1. *Doubly-charged charignos*

At the LHC, the doubly-charged charginos can be pair produced via the Drell-Yan process or associated produced with a singly-charged chargino. For light charginos around 100 GeV, the production cross section can be as large as 10 pb (It quickly drops to a few 0.1 pb for a 200 GeV $\chi^{++}$) [31]. Moreover, the singly and doubly-charged charginos decay produce multi $W-$bosons, through the typical decay topologies

$$\chi_1^+ \to \chi_1^0 W^+, \quad \chi^{++} \to \chi_1^+ W^+ \to \chi_1^0 W^+ W^+. \tag{27}$$

They are independent on the superpotential and hence are general. But note that some of $W-$bosons are off-shell, depending on the mass splitting among $\chi^{++}$ , $\chi_1^+$ and $\chi_1^0$. In the case of a sufficient degeneracy between $m_{\chi^{++}}$ and $m_{\chi_0}$, say smaller than the $W$ bosom mass threshold, both $W$ bosons are off-shell. Such a situation arises when the triplets do not significantly couple to the Higgs doublets, such as in scenario B where mixing induced splitting (or the loop correction) is only at the GeV order and moreover the LSP comes from the triplets, say in the inelastic DM scenario. Consequently the resulting leptons generically are too soft to be detected and $\chi^{++}(\chi^+)$ behavior as missing energy at the LHC (assumed to be not long-lived). The mono-jet plus missing energy signal may provide a probation to this case.

For cases with properly large mass splittings, e.g., in the scenario A, we then have a typical mass spectrum: $\chi_1^+$ with mass around 100 GeV while the heavier doubly-charged charigno mass with $m_{\chi^{++}} = \mu_T \sim 150 - 300$ GeV (see Fig. 3), as for the LSP $\chi_1^0$ having even smaller mass, e.g., 80 GeV. Consider $\chi^{\pm\pm}$ pair-production, each doubly charged charino produces missing energy and energetic same-sign leptons. Thus the tetraleptons plus missing energy signature

$$pp \to \chi^{++}\chi^{--} \to (\ell_1^+\ell_2^+)(\ell_3^-\ell_4^-)E_T^{\rm miss} \tag{28}$$

furnishes a very promising channel, of which the backgrounds are extremely suppressed, say $\sim 10^{-3}$ fb [31].

### 2. *Doubly-charged Higgs boson*

Although both scenarios present TeV scale doubly-charged Higgs bosons, such heavy particles suffer from small production rate and we thus concentrate on the clearly predicted $H^{++}$ below 100 GeV in the scenario B. Taking into account the small neutrino mass, which requires extremely tiny Yukawa couplings $(\lambda_L)_{ij} \sim \mathcal{O}(10^{-10})$, the di-boson mode $H^{++} \to W^+W^+$ thus becomes the dominant $H^{++}$ decay channel [37] and then the same-sign lepton signal search [32, 33] fails in probing/constraining this scenario. However, in addition to the small triplet VEV $v_{T_d} \sim \mathcal{O}(0.1)$ GeV suppressing, this branching decay width is suppressed by the phase space and consequently it may be long-lived at the LHC. Concretely, the width is estimated in the massless limit:

$$\Gamma(H^{++} \to W^+ f f') \sim \frac{g_2^6}{288\pi^3} \frac{v_{T_d}^2 m_{H^{++}}^3}{m_W^4}. \tag{29}$$

Taking $m_{H^{++}} = 100$ GeV and $v_{T_d} = 1$ GeV, the resulted flying distance of the decaying $H^{++}$ at the LHC is $\sim 10^{-7}$ mm. Even taking into consideration the possible large phase space suppressing, which arises when $m_{H^{++}}$ is only slightly heavier (but never lighter) than $m_W$, generically $H^{++}$ promptly decays at the collider.

Ref. [32] also considered same-sign di$-W$ search from doubly-charged Higgs, using the signature

$$pp \to \chi^{++}\chi^{--} \to (\ell_1^+\ell_2^+)(\ell_3^-\ell_4^-)E_T^{\text{miss}}, \quad \text{or} \quad (jjjj)(\ell_3^-\ell_4^-)E_T^{\text{miss}} \tag{30}$$

But it only gave a discussion on the mass region above 200 GeV. In our interesting mass region, the resulted leptons or jest tend to be soft. But its production rate is quite considerable and the pair produced $H^{++}$ can be significantly boosted, so we may use the fat-jets technique to dig out this doubly-charged Higgs boson. We leave this promising scenario for future work.

## III. DISCUSSION AND CONCLUSION

Inspired by the relatively heavy Higgs boson mass and hints for di-photon excess, and moreover guided by naturalness, we consider the low energy SUSY including light triplets with hypercharge ±1. We discuss two attractive scenarios of the TSSM:

- One is the large $\lambda_u$ and $\lambda_d$ scenario, which is able to not only enhance $m_h$ but also di-photon rate. We point out the tension between these two aspects, i.e., the latter tends to render the former excessively enhanced. But the doublet-triplets effect, favored by naturalness, can substantially reduce $m_h$ to the desired value.

- The other one is the opposite case characterized by negligibly small $\lambda_{u,d}$, which however provides an elegant explanation of the origin of the di-photon enhancement. On top of that, the model furnishes an inelastic non-thermal DM candidate, i.e., the neutral pseudo Dirac triplet fermion.

Both scenarios predict light doubly-charged objects, and thus we made a preliminary analysis their discovery prospect at the LHC, based on the signatures from same-sign $W-$bosons.

We would like to end up this paper by commenting on the grand unification prospect. When imbed the model in a $SU(5)$-GUT, the triplets with hypercharge ±1 come from symmetric rank two tensors of $SU(5)$. Under the decomposition $SU(5) \to S(3)_c \times SU(2)_L \times U(1)_Y$, we have $15 = (1,3,1) \oplus (3,2,\frac{1}{6}) \oplus (6,1,-\frac{2}{3})$ and similarly to $\overline{15}$. Adding such light particles at the weak scale will render the gauge couplings non-perturtative below the GUT-scale. For scenario A, the Landau pole is hit below the GUT-scale, and non-perturtative unification is a choice [8]. Alternatively, in scenario B, adding the magic fields can lead to gauge coupling unification as discussed in [14].

### Note added

After the completion of this work, we noticed that [34] appeared on the arxiv. This paper, working in the supersymmetric type II seesaw model and focusing on the doubly-charged Higgs boson as the source of di-photon excess as well as its LHC implication, overlaps with a part of our discussion. Our results agree with each other.

### Appendix A: Yukawa couplings' RGEs

For using in the text, here we present the RGE running of the new Yukawa couplings $\lambda_{u,d}$ as well as the running of relevant parameters, e.g., $h_t$,$h_b$ and $h_\tau$.

$$\begin{aligned}
16\pi^2\frac{d\lambda_u}{dt} &= \lambda_u\left[6h_t^\dagger h_t + 14\lambda_u^\dagger\lambda_u - 7g_2^2 - \frac{9}{5}g_1^2\right], \\
16\pi^2\frac{d\lambda_d}{dt} &= \lambda_d\left[6h_b^\dagger h_b + 2h_\tau^\dagger h_\tau + 14\lambda_d^\dagger\lambda_d - 7g_2^2 - \frac{9}{5}g_1^2\right], \\
16\pi^2\frac{dh_t}{dt} &= h_t\left[6h_t^\dagger h_t + h_b^\dagger h_b + 6\lambda_u^\dagger\lambda_u - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{13}{15}g_1^2\right], \\
16\pi^2\frac{dh_b}{dt} &= h_b\left[6h_b^\dagger h_b + h_\tau^\dagger h_\tau + h_t^\dagger h_t + 6\lambda_d^\dagger\lambda_d - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{7}{15}g_1^2\right], \\
16\pi^2\frac{dh_\tau}{dt} &= h_\tau\left[3h_b^\dagger h_b + 4h_\tau^\dagger h_\tau + 6\lambda_d^\dagger\lambda_d - 3g_2^2 - \frac{9}{5}g_1^2\right]. \qquad \text{(A1)}
\end{aligned}$$

here $t = \ln\frac{\mu}{\mu_0}$ and $\mu$ is the running scale. The presence of light triplets at the weak scale affects the MSSM gauge coupling runnings:

$$\begin{aligned}
16\pi^2\frac{dg_1}{dt} &= \frac{51}{5}g_1^3, \\
16\pi^2\frac{dg_2}{dt} &= 5g_2^3, \\
16\pi^2\frac{dg_3}{dt} &= -3g_3^3. \qquad \text{(A2)}
\end{aligned}$$

As expected, they will not show unification.

### Appendix B: The Higgs potential and mass matrix

The Higgs potential of the triplets and Higgs doublets contains three parts, among which the $F-$term and soft term can be obtained in the text, while the $D-$terms with respect to the $SU(2)_L \times U(1)_Y$ groups are given by

$$\begin{aligned}V_D =&\frac{g_1^2}{2}\left(\mathrm{Tr}(-T_u^\dagger T_u + T_d^\dagger T_d) + \frac{1}{2}\left(H_u^\dagger H_u - H_d^\dagger H_d\right)\right)^2 + \\ &\frac{g_2^2}{2}\sum_{a=1,2,3}\left[\frac{1}{2}\mathrm{Tr}(T_u^\dagger[\sigma^a, T_u]) + \frac{1}{2}\mathrm{Tr}(T_d^\dagger[\sigma^a, T_d]) + \frac{1}{2}\left(H_u^\dagger \sigma^a H_u - H_d^\dagger \sigma^a H_d\right)\right]^2 .\end{aligned} \tag{B1}$$

Collecting all the terms and adopting in the fields decomposing as done in Eq. (9), we get the CP-even Higgs mass square matrix $M_S^2$ with entries (in the basis $(h_1, h_2, h_3, h_4)$)

$$\begin{aligned}(M_S^2)_{11} \doteq& M_A^2 + m_Z^2\left(1 + \frac{\lambda_u^2 + \lambda_d^2}{g^2}\right)\sin^2 2\beta,\\ (M_S^2)_{12} \doteq& - m_Z^2\left[\cos 2\beta\left(1 + \frac{\lambda_u^2 + \lambda_d^2}{g^2}\right) - \frac{\lambda_u^2 - \lambda_d^2}{g^2}\right]\sin 2\beta,\\ (M_S^2)_{13} \doteq& m_Z\left[2\lambda_u\mu\cos 2\beta/g - (A_u + \lambda_d\mu_T)\sin 2\beta/g\right],\\ (M_S^2)_{14} \doteq& m_Z\left[2\lambda_d\mu\cos 2\beta/g + (A_d + \lambda_u\mu_T)\sin 2\beta/g\right],\\ (M_S^2)_{22} =& m_Z^2\left[\cos^2 2\beta + \frac{4}{g^2}\left(\lambda_d^2\cos^4\beta + \lambda_u^2\sin^4\beta\right)\right]\\ (M_S^2)_{23} \doteq& - 2m_Z\mathcal{M}_u/g,\\ (M_S^2)_{24} \doteq& - 2m_Z\mathcal{M}_d/g,\\ (M_S^2)_{33} \doteq& m_Z^2\mathcal{M}_u,\\ (M_S^2)_{34} \doteq& B\mu_T,\\ (M_S^2)_{44} \doteq& m_Z^2\mathcal{M}_d,\end{aligned} \tag{B2}$$

where $M_A^2 \equiv 2B\mu/\sin 2\beta$. In reality, in the heavy $m_{T_u}^2$ limit (similarly applied to $T_d^0$), $(M_S^2)_{33}$ simply assumes the form be written in a more clear way after using Eq. (6),

$$(M_S^2)_{33} \doteq m_{T_u}^2 . \tag{B3}$$

---

[1] G. Aad et al. [ATLAS Collaboration], Phys. Lett. B 716 (2012) 1; S. Chatrchyan et al. [CMS Collaboration], Phys. Lett. B 716 (2012) 30.

[2] Z. Kang, J. Li and T. Li, JHEP **1211**, 024 (2012).

[3] J. -J. Cao, et al., JHEP **1203**, 086 (2012); K. Agashe, Y. Cui and R. Franceschini, arXiv:1209.2115; S. F. King, M. Muhlleitner, R. Nevzorov and K. Walz, arXiv:1211.5074.

[4] Z. Kang, J. Li, T. Li, D. Liu and J. Shu, arXiv:1301.0453; D. G. Cerdeno, P. Ghosh and C. B. Park, arXiv:1301.1325.

[5] M. Hirsch, W. Porod, L. Reichert and F. Staub, Phys. Rev. D **86**, 093018 (2012); H. An, T. Liu and L. -T. Wang, arXiv:1207.2473; R. Huo, G. Lee, A. M. Thalapillil and C. E. M. Wagner, arXiv:1212.0560.

[6] S. P. Martin and J. D. Wells, Phys. Rev. D **86**, 035017 (2012) K. J. Bae, T. H. Jung and H. D. Kim, arXiv:1208.3748; B. Kyae and J. -C. Park, arXiv:1207.3126; M. A. Ajaib, I. Gogoladze and Q. Shafi, Phys. Rev. D **86**, 095028 (2012).

[7] T. Basak and S. Mohanty, Phys. Rev. D **86**, 075031 (2012); P. Fileviez Perez and S. Spinner, arXiv:1209.5769; I. Gogoladze, B. He and Q. Shafi, arXiv:1209.5984 [hep-ph].

[8] A. Delgado, G. Nardini and M. Quiros, arXiv:1207.6596.

[9] P. P. Giardino, K. Kannike, M. Raidal and A. Strumia, JHEP **1206**, 117 (2012); J. R. Espinosa, C. Grojean, M. Muhlleitner and M. Trott, JHEP **1212**, 045 (2012); T. Corbett, et al, Phys. Rev. D **86**, 075013 (2012); D. Carmi, A. Falkowski, E. Kuflik, T. Volansky and J. Zupan, JHEP **1210**, 196 (2012).

[10] K. Choi, S. H. Im, K. S. Jeong and M. Yamaguchi, arXiv:1211.0875; T. Gherghetta, B. von Harling, A. D. Medina and M. A. Schmidt, arXiv:1212.5243.

[11] J. R. Espinosa and M. Quiros, Phys. Lett. B **279**, 92 (1992).

[12] M. Dine, N. Seiberg and S. Thomas, Phys. Rev. D **76**, 095004 (2007).

[13] K. Agashe, A. Azatov, A. Katz and D. Kim, Phys. Rev. D **84**, 115024 (2011).

[14] P. Fileviez Perez and S. Spinner, arXiv:1211.1025 [hep-ph].

[15] C. Balazs and S. K. Gupta, arXiv:1212.1708 [hep-ph].

[16] R. N. Mohapatra and G. Senjanovic, Phys. Rev. D 23, 165 (1981); J. Schechter and J. W. F. Valle, Phys. Rev. D 22, 2227 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B 181, 287 (1981).

[17] T. Hambye, E. Ma and U. Sarkar, Nucl. Phys. B **602**, 23 (2001).

[18] M. Carena, I. Low and C. E. M. Wagner, JHEP **1208**, 060 (2012).

[19] J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, Nucl. Phys. B 106 (1976) 292; M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, Sov. J. Nucl. Phys. 30 (1979) 711 [Yad. Fiz. 30 (1979) 1368].

[20] K. Blum, R. T. D'Agnolo and J. Fan, arXiv:1206.5303 [hep-ph].

[21] Z. Kang, T. Li, T. Liu, C. Tong and J. M. Yang, Phys. Rev. D **86**, 095020 (2012).

[22] Particle Data Group Collaboration, K. Nakamura et. al., J.Phys.G G37 (2010) 075021.

[23] A. Arhrib, R. Benbrik, M. Chabab, G. Moultaka and L. Rahili, JHEP **1204**, 136 (2012); A. G. Akeroyd and S. Moretti, Phys. Rev. D **86**, 035015 (2012); L. Wang and X. -F. Han, arXiv:1209.0376; E. J. Chun, H. M. Lee and P. Sharma, JHEP **1211**, 106 (2012); W. Chao, M. Gonderinger and M. J. Ramsey-Musolf, arXiv:1210.0491; C. -S. Chen, C. -Q. Geng and L. -H. Tsai, arXiv:1212.6208; C. Han, N. Liu, L. Wu, J. M. Yang and Y. Zhang, arXiv:1212.6728.

[24] Z. Kang, T. Li, T. Liu, C. Tong and J. M. Yang, JCAP **1101**, 028 (2011).

[25] L. J. Hall, T. Moroi and H. Murayama, Phys. Lett. B **424**, 305 (1998).

[26] D. Tucker-Smith and N. Weiner, Phys. Rev. D64 (2001).

[27] V. De Romeri and M. Hirsch, JHEP **1212**, 106 (2012).

[28] E. Aprile *et al.* [XENON100 Collaboration], Phys. Rev. Lett. **109**, 181301 (2012).

[29] D. P. Finkbeiner and N. Weiner, Phys. Rev. D **76**, 083519 (2007); J. M. Cline and A. R. Frey, Annalen Phys. **524**, 579 (2012).

[30] X. Gao, Z. Kang and T. Li, arXiv:1107.3529 [hep-ph].

[31] D. A. Demir, M. Frank, K. Huitu, S. K. Rai and I. Turan, Phys. Rev. D **78**, 035013 (2008).

[32] T. Han, B. Mukhopadhyaya, Z. Si and K. Wang, Phys. Rev. D **76**, 075013 (2007).

[33] A. Melfo, M. Nemevsek, F. Nesti, G. Senjanovic and Y. Zhang, Phys. Rev. D **85**, 055018 (2012).

[34] E. J. Chun and P. Sharma, arXiv:1301.1437 [hep-ph].

[35] This is a generic conclusion if the Dirac fermions running in the loop develop a $v-$dependence only via the mixing induced by electro-weak symmetry breaking.

[36] In principle, we can keep $\lambda_u \sim \mathcal{O}(1)$ to enhance $m_h$, but it is at the price of a big fine-tuning ($\sim 1\%$) to obtain a small $v_{T_d}$.

[37] Note that the decay channel $H^{++} \to H^+W^+$ with a real $H^+$ is forbidden, since in the triplets $T_d$, the doubly-charged component is the lightest one. The reason can be explicitly found in Eq. (20), where $T_d^{++}$ receives a negative mass term from the coupling to the Higgs doublets while $T_d^+$, by contrast, gets a decrease. In other words, such a mass order is correlated with the fact that $T_d^{++}$ enhancing the di-photon rate while $T_d^+$ acts oppositely.